



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DETERMINATION OF THE AREA OF AN ELLIPSE GEOMETRICALLY, THAT OF THE CIRCLE BEING KNOWN.

BY OCTAVIAN L. MATHIOT, BALTIMORE, MD.

LET $ABDE$ represent any upright circular cylinder whose diameter is AB and perpendicular height BD . Let the cylinder be cut by a plane through D and A , the cutting edge of which is parallel to the plane of the base; then will the section $AFDH$ be an ellipse whose transverse axis AD and whose conjugate axis $FH = AB$, the diameter of the cylinder; and it is evident from the figure that this ellipse divides the cylinder into two equal parts.

Prolong the cylinder to IK so that $DI = DB$, and through IE pass a plane parallel to the plane through DA , then will the section $EMIL$ be an ellipse similar and equal to the ellipse $AFDH$.

Let the diameter of the base, $AB = b$ and the height $BD = h$, then is $AD = \sqrt{(b^2 + h^2)} = a$, say; hence a and b are the axes of the ellipse.

Produce AD to meet a perpendicular from I in O , then will the triangles ABD and IOD be similar, and we shall have $AD : AB :: ID : IO$, or, $a : b :: h : p$, where $p = IO$ the perpendicular distance between the sections through AD and EI ; hence

$$p = \frac{bh}{a}; \therefore A \frac{bh}{a} = \frac{1}{2}\pi b^2 h, \quad (1)$$

where A = the area of the ellipse;

$$\therefore A = \frac{1}{2}\pi ab. \quad (2)$$

The equality expressed by (1) is obvious from the construction of the Fig., and the result, in (2) is the recognized expression for the area of an ellipse, but, so far as I know, it has hitherto been obtained from a higher analysis than the foregoing.

In giving the above formula for the mensuration of the ellipse, a well-known author says; "this rule and the one for the following problem (the parabola) cannot be demonstrated without the aid of principles not yet considered" &c. (See Davies' Legendre, for the use of schools, p. 284.)

